



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Wild type agr-negative livestock-associated MRSA exhibits high adhesive capacity to human and porcine cells

Bünter, Julia P ; Seth-Smith, Helena M B ; Rüegg, Simon R ; Heikinheimo, Annamari ; Borel, Nicole ; Jöhler, Sophia

Abstract: Methicillin-resistant *Staphylococcus aureus* (MRSA) is one of the leading causes of nosocomial infections and a major public health concern worldwide. During the last decade, MRSA of CC398 have emerged as important colonizers of livestock. These strains also represent an increasing cause of human infections. A recent study reporting a new dominant spa type among MRSA from Finish fattening pigs (CC398/t2741) identified a strain lacking both the global virulence regulator gene locus *agr* and the adhesion gene *fnbB*. The aim of this study was to characterize this *agr/fnbB*-negative livestock-associated MRSA strain in terms of growth, hemolysis and adhesive capacity, and to provide data on its genomic background. To this end, growth curves and hemolysis patterns were generated and adhesion assays on human keratinocyte and porcine nasal mucosa cell lines were performed. Whole genome sequencing was used to determine the nature and extent of the relevant deletions in the livestock strains. For comparison, an *agr*-positive, *fnbB*-negative CC398/t2741 strain from the same pig herd, an *agr/fnbB*-positive CC398/t034 strain from another pig herd and one human MRSA strain and its isogenic Δ *agr* knockout mutant were used. The *agr*-negative strains adhered significantly better to human and porcine host cells than the *agr*-positive control strains. For the *agr*-positive porcine MRSA strains, cytotoxic effects on porcine mucosal cells were observed. The strong adhesive capacity of the naturally *agr*-negative livestock-associated MRSA, in combination with diminished cytotoxic effects, is likely favorable for inducing persistent colonization in pigs. Independently of the host cell type, similar adhesive capacities of the naturally *agr*-negative livestock-associated MRSA and the human MRSA strain were shown. Our results indicate that loss of *agr* in the livestock-associated MRSA strain investigated in this study may have increased its potential to be transmitted to and amongst humans.

DOI: <https://doi.org/10.1016/j.resmic.2016.09.006>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-127229>

Journal Article

Accepted Version

Originally published at:

Bünter, Julia P; Seth-Smith, Helena M B; Rüegg, Simon R; Heikinheimo, Annamari; Borel, Nicole; Jöhler, Sophia (2017). Wild type *agr*-negative livestock-associated MRSA exhibits high adhesive capacity to human and porcine cells. *Research in Microbiology*, 168(2):130-138.

DOI: <https://doi.org/10.1016/j.resmic.2016.09.006>

Wild type *agr*-negative livestock-associated MRSA exhibits high adhesive capacity to human and porcine cells

Julia P. Bünter¹, Helena M.B. Seth-Smith¹, Simon Rüegg², Annamari Heikinheimo⁴, Nicole Borel^{1*¶}, Sophia Johler^{3¶}

¹ Institute for Veterinary Pathology, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 268, CH-8057 Zurich, Switzerland

² Section of Epidemiology, Vetsuisse Faculty, University of Zürich, Winterthurerstrasse 270, CH-8057 Zurich, Switzerland

³ Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 272, CH-8057 Zurich, Switzerland

⁴ Department of Food Hygiene and Environmental Health, Faculty of Veterinary Medicine, University of Helsinki, Agnes Sjöberginkatu 2, P.O. Box 66, FI-00014 Helsinki, Finland

* Corresponding author

Email: n.borel@access.uzh.ch (NB)

¶These authors contributed equally to this work.

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) are one of the leading causes of nosocomial infections and a major public health concern worldwide. During the last decade, MRSA of CC398 have emerged as important colonizers of livestock. These strains also represent an increasing cause of human infections. A recent study reporting a new dominant *spa* type among MRSA from Finish fattening pigs (CC398/t2741) identified a strain lacking both the global virulence regulator gene locus *agr* and the adhesion gene *fnbB*. The aim of this study was to characterize this *agr/fnbB*-negative livestock-associated MRSA strain in terms of growth, hemolysis, and adhesive capacity, and to provide data on its genomic background. To this end, growth curves and hemolysis patterns were generated and adhesion assays on human keratinocyte and porcine nasal mucosa cell lines were performed. Whole genome sequencing was used to determine the nature and extent of the relevant deletions in the livestock strains. For comparison, an *agr*-positive, *fnbB*-negative CC398/t2741 strain from the same pig herd, an *agr/fnbB*- positive CC398/t034 strain from another pig herd, and one human MRSA strain and its isogenic Δagr knockout mutant were used. The *agr*-negative strains adhered significantly better to human and porcine host cells than the *agr*-positive control strains. For the *agr*-positive porcine MRSA strains, cytotoxic effects on porcine mucosal cells were observed. The strong adhesive capacity of the naturally *agr*-negative livestock-associated MRSA in combination with diminished cytotoxic effects is likely favorable for inducing persistent colonization in pigs. Independent of the host cell type, similar adhesive capacities of the naturally *agr*-negative livestock-associated MRSA and the human MRSA strain were shown. Our results indicate that loss of *agr* in the livestock-

46 associated MRSA strain investigated in this study may have increased its potential to be
47 transmitted to and amongst humans.

48 Introduction

49

50 Livestock-associated MRSA (LA-MRSA) strains have emerged worldwide. The strains
51 typically belong to CC398 and are detected in particularly high rates among pigs [1-4]. LA-
52 MRSA CC398 are thought to have originated from human methicillin-susceptible
53 *Staphylococcus aureus* and acquired methicillin resistance during the host jump, at the same
54 time losing genes important for colonization and infection in humans [5]. Even though LA-
55 MRSA of CC398 lack important virulence factors, they have been suggested to exhibit an
56 enhanced ability to acquire virulence factors through mobile genetic elements [6].

57 Transmission of CC398 from animals to humans has been described and occupational
58 exposure to pigs was shown to increase nasal carriage rates (pig farmers: 20-86%;
59 veterinarians: 4.6-45%) compared to the general population (0.8-3%) [2, 4, 7-14]. In addition,
60 in recent years, there have been an increasing number of reports of MRSA CC398 strains
61 causing invasive infections in humans without contact to livestock [15-18]. Therefore, the
62 transmissibility and the virulence potential of LA-MRSA of CC398 need to be closely
63 monitored.

64 Heikinheimo et al. [19] recently identified a new dominant *spa* type among LA-MRSA
65 strains colonizing Finnish fattening pigs (CC398/t2741). Interestingly, for one CC398/t2741
66 strain isolated in this study, no genes within the accessory gene regulator locus (*agr*) or the
67 fibronectin-binding protein B (*fnbB*) were detected by DNA microarray. The quorum sensing
68 *agr* system is a major virulence gene regulator. Activation of *agr* has been shown to inhibit
69 expression of certain cell-wall associated proteins and to increase exoprotein expression
70 during the post-exponential phase of growth [20, 21]. While *agr* plays a major role in acute

infections, *agr*-defective mutants can frequently be isolated from bacteraemic patients and were linked to persistent infections [22] and increased mortality [23]. The *fnbB* gene encodes the fibronectin binding protein B (FnBPB), which belongs to the group of microbial surface components recognizing adhesive matrix molecules (MSCRAMMs). It facilitates adhesion of *S. aureus* to the components of the host cells and has a role in invasion [24]. Combined FnBPA and FnBPB function has been shown to be essential to induce severe infection [25]. The aim of this study was to characterize the naturally *agr/fnbB*-negative CC398/t2741 LA-MRSA strain detected in Finland with regard to growth, hemolysis, and adhesive capacity, and to describe the genomic background of the strain.

80

81 **Methods**

82 **Bacterial strains**

An overview of the strains used in this study is presented in Table 1. The naturally *agr/fnbB*-negative LA-MRSA strain investigated in this study (Fin47_H17) was isolated from the carcass of a fattening pig in Finland [19]. Two porcine LA-MRSA strains isolated in the same study were used as controls: Fin46_H17, an *agr*-positive, *fnbB*-negative isolate from the same herd (herd 17), and Fin48_H18, an *agr/fnbB*-positive isolate from another herd (herd 18). As additional controls, a MRSA strain that had caused recurring skin infections in humans (MN10) [26] and its isogenic *agr* knockout mutant (MN10_ Δ *agr*) were used. MN10_ Δ *agr* was obtained by transduction of the *agr* knockout from RN6911 to MN10 using phage 80 α and protocols previously described [27]. Correct deletion of *agr* in putative mutants grown on tetracycline selective plates was confirmed by PCR.

93

94 **Table 1: Strains used in this study.** Presence/absence of *agr* and *fnbB* is indicated as
 95 previously determined by DNA microarray [19].

| Strain ID | Characteristics | Country | Year | Source [reference] |
|-------------------------|---|---------|------|----------------------------|
| Fin47_H17 | CC398/t2741, <i>agr</i> ⁻ , <i>fnbB</i> ⁻ | FI | 2015 | Pig carcass, herd 17 [19] |
| Fin46_H17 | CC398/t2741, <i>agr</i> ⁺ , <i>fnbB</i> ⁻ | FI | 2015 | Pig nares, herd 17 [19] |
| Fin48_H18 | CC398/t034, <i>agr</i> ⁺ , <i>fnbB</i> ⁺ | FI | 2015 | Pig nares, herd 18 [19] |
| MN10 | CC5/t002, <i>agr</i> ⁺ , <i>fnbB</i> ⁺ | CH | 2013 | Human skin infection [26] |
| MN10_Δ<i>agr</i> | CC5/t002, NM10 <i>agr</i> :: <i>tetM</i> , <i>agr</i> ⁻ , <i>fnbB</i> ⁺ | CH | 2013 | This study |
| RN27 | 80alpha lysogen | - | - | Brigitte Berger-Bächi [28] |
| RN6911 | <i>agr</i> operon deleted (<i>agr</i> :: <i>tetM</i>) | - | - | Brigitte Berger-Bächi [28] |

96

97 **Growth parameters**

98 Single colonies were transferred from 5% sheep blood agar to 50 mL of Luria Bertani
99 (LB) broth (Becton Dickinson, Allschwil, Switzerland) and grown for 24 h at 37°C (225
100 rpm). Growth of all strains in LB was determined by viable cell counts after 3 h, 6 h, 9 h, and
101 24 h using 10-fold dilution series and plate count agar (Sigma-Aldrich, Stockholm, Sweden),
102 with incubation at 37°C for 18-24 h. Growth parameters such as exponential phase growth
103 rate and maximum cell density were determined using DMFit 3.0 [29].

105 **Hemolysis**

106 Screening for hemolytic activity was performed as previously described [30]. Briefly,
107 alpha- and delta-hemolytic activity was determined by perpendicular streaking to the beta-
108 hemolysin producing *S. aureus* reference strain RN4220 on 5% sheep blood agar and
109 incubation over night at 37°C. In this assay, beta hemolysis results in a turbid zone. Alpha
110 hemolytic activity of the test strain is inhibited by beta hemolysis of RN4220 where the
111 strains intersect. Delta hemolysis of the test strain is synergistic with beta hemolysin of
112 RN4220, resulting in an amplified zone of clearing where the strains intersect [31]. Gamma
113 hemolytic activity cannot be detected in this assay, as it is inhibited by agar [32]. Subsequent
114 cold shock (exposure to 4°C for 12h) was used to determine beta-hemolysis activity.

116 **Adhesion assay**

117 Two different cell lines were used for adhesion assays: human cell line HaCaT
118 (Human adult low Calcium high Temperature keratinocytes, CLS Cell Lines Service GmbH,

119 Eppelheim, Germany; [33]) and porcine cell line PT-K75 (porcine nasal turbinate/mucosa,
120 CRL-2528, ATCC, Manassas, USA). The medium for the adhesion assay consisted of
121 DMEM supplemented with 4.5 g/L glucose, 584 mg/L L-glutamine and 10% FCS (CLS Cell
122 Lines Service GmbH) for the HaCaT cells and DMEM supplemented with 4.5 g/L glucose,
123 580mg/L L-glutamine, 110mg/L sodium pyruvate (GIBCO, Thermo Fisher Scientific,
124 Invitrogen, Carlsbad, CA, USA) and 10% FCS (BioConcept, Allschwil, Switzerland) for the
125 PT-K75 cells.

126 After at least one passage in a 75cm² canted neck tissue culture flask with vented cap
127 (Corning, Sigma-Aldrich), HaCaT-cells were seeded at a concentration of 2×10^5 cells/well
128 and PT-K75-cells at a concentration of 1.25×10^5 cells/well in a 24-well flat-bottom cell
129 culture plate with low-evaporation lid (Techno Plastic Products AG, TPP, Trasadingen,
130 Switzerland). The culture plates were incubated for 24 hours at 37°C and 5% CO₂ to reach a
131 confluency of at least 80%.

132 Bacterial strains were prepared by growth on 5% sheep blood agar overnight at 37°C.
133 Overnight cultures were prepared by inoculation of 5 mL LB broth with a single colony and
134 incubation at 37°C (225 rpm shaking) for 15 hours. Cultures were subsequently adjusted to
135 OD₅₉₀ = 0.40 ($\sim 10^8$ CFU/ml) for each strain. The bacteria were used at a multiplicity of
136 infection (MOI) of 10 to infect both cell lines (HaCaT and PT-K75) in 24-well plates.
137 Uninfected cells were used as negative controls. After 15 min of incubation at 37°C, cells
138 were imaged, before monolayers were washed eight times with Dulbecco's phosphate
139 buffered saline (DPBS, GIBCO) and cells were harvested after five minutes of incubation
140 with Trypsin-EDTA (0.25%, GIBCO) by scraping with a cell scraper (TPP). Bacterial cell
141 counts were determined by 10-fold dilution series and plate count agar. Three independent

142 biological experiments were performed, each using three replicates per strain. For imaging
143 purposes, the Nikon Eclipse Ti-U inverted microscope was used at 200-fold magnification
144 with software NIS-Elements AR Analysis 4.3 (Nikon AG, Egg, Switzerland).

145

146 **Statistical analysis**

147 All statistical analyses with the exception of growth parameter analyses were
148 performed using R, 3.2. Results were compared using a gaussian linear model of the form
149 $\text{count} \sim \text{strain} + \text{host.cellline}$. Growth parameters were compared between strains using SPSS
150 Statistics 22 (SPSS Inc.Chicago, Illinois). Growth parameters such as exponential phase
151 growth rate and maximum cell density were determined using DMFit 3.0 (Baranyi &
152 Roberts, 1994) and compared using one-way ANOVA. Differences were considered
153 statistically significant if $p < 0.05$.

154

155 **Whole genome sequencing**

156 DNA was extracted from overnight cultures grown on 5% sheep blood agar plates
157 using the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany) according to the
158 manufacturer's instructions. The porcine LA-MRSA strains Fin47_H17, Fin46_H17, and
159 Fin48_H18 were sequenced on the Illumina Miseq platform with 250bp paired end reads
160 following NEBNextUltra library creation. Raw sequencing reads were adaptor and quality
161 trimmed and filtered using Trimmomatic version 0.32 [34]. Mapping against the CC398
162 reference strain S0385 (EMBL Accession AM990992) [6] using BWA [35] indicated
163 chromosome coverage of each genome to be: Fin47_H17 = $132.8x \pm 76.2$, Fin46_H17 =

164 140.6 x \pm 71.2, Fin48_H18 = 115.1x \pm 53.7. *De novo* assembly used SPAdes v 3.1.0 in multi-
165 cell mode [36]. Scaffolds were rearranged in ACT [37] against S0385 using files generated in
166 WebACT (<http://www.webact.org/>). Resulting assemblies comprise 28, 25 and 27 scaffolds
167 respectively, with scaffolds under 1kb, which did not match the reference, excluded. Target
168 coding sequences (CDSs) involved in adhesion were identified and analysed in all genomes
169 using ACT and, where necessary, single CDSs were aligned using ClustalW. The phylogeny
170 was generated in <http://wgsastaging.pathogensurveillance.net/> using available assembled
171 CC398 genomes for context [6, 38-42]. Antibiotic resistance phenotypes were also predicted
172 using this online tool.
173 All read data and assemblies have been deposited with ENA under project number
174 (PRJEB14187).

175

Results

Growth parameters

No statistically significant differences were observed between growth of the five strains in LB broth with regard to exponential phase growth rate or maximum cell density (Fig. 1). However, the naturally *agr*-negative LA-MRSA Fin47_H17 exhibited the lowest exponential phase growth rate ($\mu_{\max} = 0.50$) compared to all other strains ($\mu_{\max} = 0.51$ - 0.57). Fin47_H17 also reached the lowest maximum cell density ($8.97 \log \text{ CfU/mL}$) compared to all other strains tested (9.70 - $9.91 \log \text{ CfU/mL}$).

Fig. 1: Growth of all tested strains in LB over 24h. The tested strains exhibited no statistically significant differences in exponential phase growth rate or maximum cell density reached (error bars: 95% CI).

Haemolysis pattern

Hemolysis patterns after incubation at 37°C overnight and after subsequent cold exposure are shown in Fig. 2. No hemolysis was visible for both the naturally *agr*-negative Fin47_H17 strain and *agr* knockout mutant MN10_ Δagr . Parental strain MN10 exhibited alpha and delta hemolysis. Fin46_H17 exhibited alpha and beta hemolysis, and Fin48_H18 exhibited beta and delta hemolysis.

Fig. 2: Hemolysis. The *agr*-negative strains exhibited no hemolysis (Fin47_H17 and MN_Δ*agr*). MN10 exhibited alpha and delta hemolysis, Fin46_H17 exhibited alpha and beta hemolysis, and Fin48_H18 exhibited beta and delta hemolysis.

Adhesion properties

While no morphological changes were observed in HaCaT cells after bacterial infection, infection of PT-K75 cells with the *agr*-positive porcine LA-MRSA strains (Fin46_H17 and Fin48_H18) resulted in a cytotoxic effect including shrinking, rounding and detachment of the cells (Fig. 3). To minimize cytotoxicity, a trial experiment was performed on PT-K75 cells with an MOI of 1: cytotoxic effects were still observed (data not shown). In contrast, neither the naturally *agr*-negative porcine LA-MRSA strain Fin47_H17 nor the human-associated MRSA strain MN10 and its isogenic *agr* knockout mutant induced any morphological changes in the PT-K75 cells (Fig. 3).

Counts of adherent bacteria revealed that the *agr*-negative strains Fin47_H17 and MN10_Δ*agr* displayed increased adhesive capacity in both human HaCaT and porcine PT-K75 cells (Fig. 4) compared to the *agr*-positive LA-MRSA and the human-associated parental MRSA strain MN10, respectively. Median colony counts for Fin47_H17 were 3,600 for HaCaT and 14,000 for PT-K75 cells per culture well. These results show significantly higher adhesive capacity of this strain compared to the *agr*-positive strains Fin46_H17 (790 to HaCaT, $p = 1.3e-06$), Fin48_H18 (500 to HaCaT, $p = 2.0e-07$) or MN10 (1,200 to HaCaT, $p = 4.3e-13$; 7,200 to PT-K75, $p = 0.001$). The median colony counts for the MN10_Δ*agr* mutant strain (3,190 to HaCaT and 35,000 to PT-K75) were significantly higher than those of MN10 (1,200 to HaCaT, $p = 0.003$; 7,200 to PT-K75, $p = 4.5e-05$). Thus, the naturally *agr*-

219 negative LA-MRSA and MN10_Δagr exhibited significantly higher adhesive capacity than
220 the agr-positive LA-MRSA strains and parental strain MN10, respectively. Overall, the tested
221 porcine and human MRSA strains adhered significantly better to the porcine than to the
222 human cells ($p = 1.3e-06$).

223
224 **Fig. 3: Cytotoxic effects on porcine cells.** Representative images of the two cell lines before
225 (T_0) and shortly after ($T_{15 \text{ min}}$) infection. Bacterial strains used were the porcine LA-MRSA
226 strains Fin47_H17, Fin46_H17, and Fin48_H18, as well as the MRSA strain linked to human
227 infections (MN10) and its isogenic knockout mutant MN10_Δagr. Cytotoxic effects of the
228 agr-positive porcine LA-MRSA strains Fin46_H17 and Fin48_H18 on PT-K75 cells can be
229 identified at $T_{15 \text{ min}}$.

230
231 **Fig. 4: Adhesion to A) human cells and B) porcine cells: colony counts for different**
232 **strains.** The plots indicate the median (horizontal line), the range from the first to the third
233 quartile (25–75%; box), and the extreme values (whiskers) of the colony counts for each
234 strain after adhesion to A) human cells (HaCaT) and B) porcine cells (PT-K75). The data
235 resulted from three independent biological experiments with three replicates for each strain
236 ($n=9$).

237

238 **Genomic analysis**

239 Whole genome sequencing and phylogenetic analysis of strains within CC398
240 confirmed that Fin47_H17 is closely related to Fin46_H17, with Fin48_H18 more distantly
241 related (S1 Fig).

242 In the naturally *agr*-negative LA-MRSA Fin47_H17, a deletion of 3,789 bp covering *hld*,
243 *agrB* and *agrC* (Fig. 5A) was identified. In both this strain and the closely related strain
244 Fin46_H17, a further deletion completely removing *fnbB* was detected that leaves the
245 adjacent *fnbA* intact (Fig. 5B). However, as *fnbB* in both Fin48_H18 and reference strain
246 S0385 is truncated after 349 amino acids with a single nucleotide mutation causing a
247 premature stop codon, the functional relevance of this is unclear.

248 Genes associated with adhesion were investigated within the genomes (S2 Table). Such genes
249 often have varying repeat lengths, and this was identified in these strains in the genes: *spa*,
250 *sdrC* and the hypothetical protein encoding CDS *SAPIG1791*; with the further genes *coa*,
251 *sdrD*, *clfA*, *clfB*, *ebhA* and *cna* almost identical between the three porcine LA-MRSA strains,
252 but variable in comparison to homologues in S0385. The *vwb* gene, encoding a secreted von
253 Willebrand factor-binding protein, was found to be absent at the equivalent chromosomal
254 locus in Fin48_H18, with a 13.7 kb deletion covering this region. All other genes investigated
255 were found to be identical or almost identical to those in S0385 (*eno*, *isdB*, *isdA*, *ecb*, *fib*,
256 *ebpS*, *map*, *sdrH*, *agrA*, *isaB*).

257 The strains carry alternative complements of plasmids and phage (S2 Table) and the
258 predicted antibiotic resistance profiles (Fin47_H17 and Fin46_H17: PEN^R, TET^R, ERY^R,

259 MET^R; Fin 48_H18: PEN^R, TET^R, MET^R, TRI^R) largely correspond to those phenotypically
260 determined previously, with the exception of clindamycin [19].

261

262 **Fig. 5: Comparison of genome loci around *agr* and *fnbB*.** Each line refers to a genomic
263 section in the strain indicated. Arrows indicate CDSs. Grey bars show homology between
264 loci according to the scale. **A.** *agr* locus showing the extent of the deletion in Fin47_H18,
265 covering *SAPIG2071-2075*. *hld* is shown in yellow and the *agr* locus (*agrBCA*) in green.
266 Only *argA* remains in Fin47_H17. **B.** *fnbB* locus showing the deletion of *fnbB* (*SAPIG2551*)
267 in both Fin47_H17 and Fin46_H17. *fnbB* in both S0385 and Fin48_H18 is truncated by a
268 stop codon and is shown in brown for a pseudogene. *fnbA* (yellow) displays some homology
269 to *fnbB*. Figure was drawn in Easyfig [43].

270

271 Discussion

272 Heikinheimo et al. [19] investigated LA-MRSA isolated from Finnish fattening pigs
273 at slaughter. They found that CC398/t2741 strains were predominant among the isolated LA-
274 MRSA. Interestingly, one of the isolated strains (Fin47_H17) did not harbor genes from the
275 *agr* locus. In this study, we determined the genomic basis of this loss of *agr* and its effect on
276 growth, hemolytic activity, and adhesive capacity to human and porcine cell lines.

277 Loss of *agr* did not result in altered growth parameters, but had a strong impact on
278 hemolytic activity consistent with previous studies suggesting that expression of alpha and
279 delta hemolysins is strongly induced by *agr* [30, 31]. Hemolysins are widely recognized as

280 important virulence factors in *S. aureus* infections. However, selective survival of *agr*-
281 defective non-hemolytic *S. aureus* in wound and abscess models has been demonstrated [44].

282 In this study, both *agr*-negative strains tested exhibited increased adhesive capacity to
283 both human and porcine cells. These results are consistent with previous findings suggesting
284 *agr* to inhibit the expression of adhesion proteins [45-48]. Loss of *agr* was shown to
285 significantly increase adhesive capacity to human endothelial [49] and mesothelial cells [50].
286 In a bovine mammary epithelial cell line (MAC-T), two- to threefold higher numbers of
287 internalized viable bacteria of an *agr* mutant were recovered compared to the wild type [51].

288 LA-MRSA CC398 was reported to cause more cell damage within epithelial cells
289 than community-associated and hospital-acquired MRSA [52]. In this study, pronounced
290 cytotoxic effects of the *agr*-positive porcine LA-MRSA strains on porcine cells were
291 identified. In contrast, the closely related *agr*-negative porcine LA-MRSA Fin47_H17 did not
292 induce cytotoxic effects. This may be due to reduced expression levels of *agr*-dependent
293 cytotoxic proteins such as alpha hemolysin [53]. It is also consistent with a previous study
294 reporting cytotoxic effects in MAC-T cells upon infection with RN6390, but no cytotoxic
295 effects in *agr* mutant strain RN6911 [51]. The finding that porcine *agr*-positive LA-MRSA
296 only led to cytotoxicity in the porcine cell line could be due to host-specific factors
297 influencing susceptibility to alpha hemolysin [54].

298 Whole genome sequencing revealed that the *fnbB* gene in Fin48_H18 is truncated, potentially
299 impairing functionality of FnBPB, whereas it is fully deleted from the genomes of strains
300 Fin47_17 and Fin46_17. We observed no significant differences concerning adherence to
301 porcine cells of Fin48_H18 and Fin46_17. For strain Newman, it has been suggested that
302 truncation of *fnbB* was transferred to *fnbA*, resulting in complete secretion FnBPs to the

culture medium and loss of cell wall anchor function [55]. Consistent with these findings, reduced adhesive capacity of a *fnbAB* mutant to HaCaT cells has been described [56].

We observed no significant differences in adhesive capacity between porcine and human MRSA apart from the described influence of *agr*. However, irrespective of strain origin (LA-MRSA or human MRSA), adherence to porcine cells significantly exceeded adherence to human cells. LA-MRSA CC398 do not only colonize livestock, but can also be transmitted to humans with direct livestock exposure or even between humans. Although decreased adherence of LA-MRSA compared to community/hospital-associated MRSA to human endothelial and epithelial cells was observed [52], LA-MRSA strains may rapidly adapt to the human host. Our results suggest that LA-MRSA are capable of adherence to both porcine and human cells with the same efficiency as a community-acquired MRSA control strain linked to recurring infections within a family [26].

Conclusion

We were able to show that the recently reported naturally *agr*-negative and *fnbB*-negative LA-MRSA strain Fin47_H17 exhibits high adhesive capacity to porcine and human cells. Our results suggest that loss of *agr* leads to increased adhesive capacity and that porcine LA-MRSA can exhibit pronounced *agr*-dependent cytotoxic effects on porcine cells. Our findings provide further evidence for the importance of LA-MRSA as an emerging public health concern.

Acknowledgments

The authors are grateful to Prof. Roger Stephan, Prof. Angelika Lehner, Dr. Taurai Tasara, Dr. Claudia Guldemann and Dr. Henna-Maria Sihto from the Institute for Food Safety and Hygiene, University of Zurich and Dr. Cory Ann Leonard, Dr. Hanna Marti, med. vet. Sabrina Wanninger and med. vet. Jasmin Kuratli from the Institute for Veterinary Pathology, University of Zurich, Vetsuisse Faculty, for helpful advice, discussion and laboratory support. We also thank the Functional Genomics Centre Zurich for technical assistance.

Conflict of interest statement

The authors declare that they have no conflict of interest with respect to the research, authorship, and/or publication of this article.

References

- Alt K, Fetsch A, Schroeter A, Guerra B, Hammerl JA, Hertwig S, et al. Factors associated with the occurrence of MRSA CC398 in herds of fattening pigs in Germany. *BMC Vet Res*. 2011;7:69. doi: 10.1186/1746-6148-7-69.
- Smith TC, Male MJ, Harper AL, Kroeger JS, Tinkler GP, Moritz ED, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) strain ST398 is present in midwestern U.S. swine and swine workers. *PLoS One*. 2009;4(1):e4258. doi: 10.1371/journal.pone.0004258.

345

346 de Neeling AJ, van den Broek MJ, Spalburg EC, van Santen-Verheuvél MG, Dam-Deisz

347 WD, Boshuizen HC, et al. High prevalence of methicillin resistant *Staphylococcus aureus* in

348 pigs. Vet Microbiol. 2007;122(3-4):366-72

349

350 Khanna T, Friendship R, Dewey C, Weese JS. Methicillin resistant *Staphylococcus aureus*

351 colonization in pigs and pig farmers. Vet Microbiol. 2008;128(3-4):298-303.

352

353 Price LB1, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. *Staphylococcus*

354 *aureus* CC398: host adaptation and emergence of methicillin resistance in livestock. MBio.

355 2012;3(1). pii: e00305-11. doi: 10.1128/mBio.00305-11. Print 2012.

356

357 Schijffelen MJ, Boel CH, van Strijp JA, Fluit AC 2010. Whole genome analysis of a

358 livestock-associated methicillin-resistant *Staphylococcus aureus* ST398 isolate from a case of

359 human endocarditis. BMC Genomics 2010;11: 376. doi: 10.1186/1471-2164-11-376

360

361 Ellis MW, Hospenthal DR, Dooley DP, Gray PJ, Murray CK. Natural history of community-

362 acquired methicillin-resistant *Staphylococcus aureus* colonization and infection in soldiers.

363 Clin Infect Dis. 2004;39(7):971-9.

364 Kuehnert MJ, Kruszon-Moran D, Hill HA, McQuillan G, McAllister SK, Fosheim G, et al.

365 Prevalence of *Staphylococcus aureus* nasal colonization in the United States, 2001-2002. J

366 Infect Dis. 2006;193(2):172-9.

367

368 Wulf M, van Nes A, Eikelenboom-Boskamp A, de Vries J, Melchers W, Klaassen C, et al.
 369 Methicillin-resistant *Staphylococcus aureus* in veterinary doctors and students, the
 370 Netherlands. Emerg Infect Dis. 2006;12(12):1939-41.
 371
 372 Wulf MW, Sørnum M, van Nes A, Skov R, Melchers WJ, Klaassen CH, et al. Prevalence of
 373 methicillin-resistant *Staphylococcus aureus* among veterinarians: an international study. Clin
 374 Microbiol Infect. 2008;14(1):29-34.
 375
 376 van Belkum A, Melles DC, Peeters JK, van Leeuwen WB, van Duijkeren E, Huijsdens XW,
 377 et al. Methicillin-resistant and -susceptible *Staphylococcus aureus* sequence type 398 in pigs
 378 and humans. Emerg Infect Dis. 2008;14(3):479-83. doi: 10.3201/eid1403.070760.
 379
 380 Cuny C, Nathaus R, Layer F, Strommenger B, Altmann D, Witte W. Nasal colonization of
 381 humans with methicillin-resistant *Staphylococcus aureus* (MRSA) CC398 with and without
 382 exposure to pigs. PLoS One. 2009;4(8):e6800. doi: 10.1371
 383
 384 Denis O, Suetens C, Hallin M, Catry B, Ramboer I, Dispas M, et al. Methicillin-resistant
 385 *Staphylococcus aureus* ST398 in swine farm personnel, Belgium. Emerg Infect Dis.
 386 2009;15(7):1098-101. doi: 10.3201/eid1507.080652.
 387
 388 Huber H, Giezendanner N, Stephan R, Zweifel C. Genotypes, Antibiotic Resistance Profiles
 389 and Microarray-Based Characterization of Methicillin-Resistant *Staphylococcus aureus*
 390 Strains Isolated from Livestock and Veterinarians in Switzerland. Zoonoses Public Health.
 391 2011;58(5):343-9. doi: 10.1111/j.1863-2378.2010.01353.x.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

392

393 Wulf MW, Markestein A, van der Linden FT, Voss A, Klaassen C, Verduin CM. First
 394 outbreak of methicillin-resistant *Staphylococcus aureus* ST398 in a Dutch hospital, June
 395 2007. Euro Surveill. 2008;13(9). pii: 8051.

396

397 Valentin-Domelier AS, Girard M, Bertrand X, Violette J, François P, Donnio P-Y, et al.
 398 Methicillin-susceptible ST398 *Staphylococcus aureus* responsible for bloodstream infections:
 399 an emerging human-adapted subclone? PLoS One. 2011;6:e28369.
 400 10.1371/journal.pone.0028369

401

402 Uhlemann AC, Porcella SF, Trivedi S, Sullivan SB, Hafer C, Kennedy AD, et al.
 403 Identification of a highly transmissible animal-independent *Staphylococcus aureus* ST398
 404 clone with distinct genomic and cell adhesion properties. MBio. 2012;3(2). pii: e00027-12.
 405 doi: 10.1128/mBio.00027-12.

406

407 Verkade E, Bergmans AM, Budding AE, van Belkum A, Savelkoul P, Buiting AG, et al.
 408 Recent emergence of *Staphylococcus aureus* clonal complex 398 in human blood culture.
 409 PLoS ONE 2012;7:e41855.

410

411 Heikinheimo A, Johler S, Karvonen L, Julmi J, Fredriksson-Ahomaa M, Stephan R. New
 412 dominant spa type t2741 in livestock-associated MRSA (CC398-MRSA-V) in Finnish
 413 fattening pigs at slaughter. Antimicrob Resist Infect Control. 2016;5:6. doi: 10.1186/s13756-
 414 016-0105-8.

416 Bronner S, Monteil H, Prevost G. Regulation of virulence determinants in *Staphylococcus*
 417 *aureus*: complexity and applications. FEMS Microbiol. Rev. 2004;28:183–200.
 418
 419 Tseng CW, Stewart GC. Rot repression of enterotoxin b expression in *Staphylococcus aureus*.
 420 J. bacteriol., 2005 July 21. ed. 187:5301–5309.
 421
 422 Fowler VGJr, Sakoulas G, McIntyre LM, Meka VG, Arbeit RD, Cabell CH, et al. Persistent
 423 bacteremia due to methicillin-resistant *Staphylococcus aureus* infection is associated with agr
 424 dysfunction and low-level in vitro resistance to thrombin-induced platelet microbicidal
 425 protein. J Infect Dis. 2004;190:1140–9.
 426
 427 Schweizer ML, Furuno JP, Sakoulas G, Johnson JK, Harris AD, Shardell MD, et al.
 428 Increased mortality with accessory gene regulator (agr) dysfunction in *Staphylococcus aureus*
 429 among bacteremic patients. Antimicrob Agents Chemother. 2011;55:1082–7.
 430
 431 Foster TJ, Geoghegan JA, Ganesh VK, Höök M. Adhesion, invasion and evasion: the many
 432 functions of the surface proteins of *Staphylococcus aureus*. Nat Rev Microbiol. 2013;12:49–
 433 62.
 434
 435 Shinji H, Yosizawa Y, Tajima A, Iwase T, Sugimoto S, Seki K, et al. Role of fibronectin-
 436 binding proteins A and B in in vitro cellular infections and in vivo septic infections by
 437 *Staphylococcus aureus*. Infect Immun. 2011;79:2215–23.
 438

439 Nüesch-Inderbinnen MT, Stalder U, Johler S, Hächler H, Stephan R, Nüesch H-J. Intrafamilial
440 spread of a Panton-Valentine leukocidin-positive community-acquired methicillin-resistant
441 *Staphylococcus aureus* belonging to the paediatric clone ST5 SSCmecIV. JMM Case Reports
442 2014 Sept 12. doi:10.1099/jmmcr.0.001859.

443
444 Charpentier E, Anton AI, Barry P, Alfonso B, Fang Y, Novick RP. Novel cassette-based
445 shuttle vector system for gram-positive bacteria. Appl Environ Microb. 2004;70:6076–85.

446
447 Novick RP. Analysis by transduction of mutations affecting penicillinase formation in
448 *Staphylococcus aureus*. J Gen Microbiol. 1963;33:121–36.

449
450 Baranyi J, Roberts TA. A dynamic approach to predicting bacterial growth in food. Int J Food
451 Microbiol. 1994;23(3-4):277-94.

452
453 Herbert S, Ziebandt AK, Ohlsen K, Schäfer T, Hecker M, Albrecht D, et al. Repair of global
454 regulators in *Staphylococcus aureus* 8325 and comparative analysis with other clinical
455 isolates. Infect Immun. 2010;78(6):2877-89. doi: 10.1128/IAI.00088-10.

456
457 Adhikari RP, Arvidson S, Novick RP. A nonsense mutation in agrA accounts for the defect in
458 agr expression and the avirulence of *Staphylococcus aureus* 8325-4 traP::kan. Infect. Immun.
459 2007;75:4534–4540.

460
461 Wiseman, GM. The hemolysins of *Staphylococcus aureus*. Bacteriol. Rev. 1975;39:317–344.

462

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

463 Boukamp P, Petrussevska RT, Breitkreutz D, Hornung J, Markham A, Fusenig NE. Normal
 464 keratinization in a spontaneously immortalized aneuploid human keratinocyte cell line. *J Cell*
 465 *Biol.* 1988;106(3):761-71.
 466
 467 Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
 468 data. *Bioinformatics* 2014;30:2114-2120. doi: 10.1093/bioinformatics/btu170
 469
 470 Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler Transform.
 471 *Bioinformatics* 2009;25:1754-1760.
 472
 473 Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a
 474 new genome assembly algorithm and its applications to single-cell sequencing. *J Comput*
 475 *Biol* 2012;19: 455-477. doi: 10.1089/cmb.2012.0021.
 476
 477 Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. ACT: the
 478 Artemis Comparison Tool. *Bioinformatics* 2005;21:3422-3423. doi:
 479 10.1093/bioinformatics/bti553
 480
 481 Uhlemann AC, Porcella SF, Trivedi S, Sullivan SB, Hafer C, Kennedy AD, et al.
 482 Identification of a highly transmissible animal-independent *Staphylococcus aureus* ST398
 483 clone with distinct genomic and cell adhesion properties. *MBio.* 2012 Feb 28;3(2). pii:
 484 e00027-12. doi: 10.1128/mBio.00027-12. Print 2012.
 485

486 van der Mee-Marquet N, Hernandez D, Bertrand X, Quentin R, Corvaglia AR, François P.
 487 Whole-Genome Sequence of the Ancestral Animal-Borne ST398 *Staphylococcus aureus*
 488 Strain S123. Genome Announc. 2013 Aug 29;1(5). pii: e00692-13. doi:
 489 10.1128/genomeA.00692-13.
 490
 491 Hernandez D, van der Mee-Marquet N, Kluytmans J, Donnio PY, Quentin R, Corvaglia AR,
 492 et al. Whole-Genome Sequences of *Staphylococcus aureus* ST398 Strains of Animal Origin.
 493 Genome Announc. 2013 Sep 26;1(5). pii: e00689-13. doi: 10.1128/genomeA.00689-13.
 494
 495 Corvaglia AR1, François P, Bertrand X, Quentin R, Hernandez D, van der Mee-Marquet N.
 496 Whole-Genome Sequences of Two *Staphylococcus aureus* ST398 Strains of Human Origin,
 497 S94 and S100. Genome Announc. 2013 Aug 29;1(5). pii: e00691-13. doi:
 498 10.1128/genomeA.00691-13.
 499
 500 Golding GR, Bryden L, Levett PN, McDonald RR, Wong A, Graham MR, et al. Whole-
 501 genome sequence of livestock-associated st398 methicillin-resistant *staphylococcus aureus*
 502 Isolated from Humans in Canada. J Bacteriol. 2012 Dec;194(23):6627-8. doi:
 503 10.1128/JB.01680-12.
 504
 505 Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
 506 Bioinformatics. 2011 Apr 1;27(7):1009-10. doi: 10.1093/bioinformatics/btr039.
 507

508 Schwan WR, Langhorne MH, Ritchie HD, Stover CK. Loss of hemolysin expression in
 509 *Staphylococcus aureus* agr mutants correlates with selective survival during mixed infections
 510 in murina abscesses and wounds. FEMS Immunol Med Microbiol. 2003;38(1):23-8.
 511
 512 Patel AH, Kornblum J, Kreiswirth B, Novick R, Foster TJ. Regulation of the protein A-
 513 encoding gene in *Staphylococcus aureus*. Gene. 1992;114:25–34
 514
 515 Saravia-Otten P, Müller HP, Arvidson S. Transcription of *Staphylococcus aureus* fibronectin
 516 binding protein genes is negatively regulated by agr and an agr-independent mechanism. J.
 517 Bacteriol. 1997;179:5259–5263.
 518
 519 Projan SJ, Novick RP. The molecular basis of pathogenicity. In: Crossley KB, Archer GL,
 520 editors. The staphylococci in human disease. Churchill Livingstone, New York, N.Y; 1997.
 521 pp 55–82.
 522
 523 Wolz C, Pöhlmann-Dietze P, Steinhuber A, Chien YT, Manna A, van Wamel W, et al. Agr-
 524 independent regulation of fibronectin-binding protein(s) by the regulatory locus sar in
 525 *Staphylococcus aureus*. Mol Microbiol. 2000;36(1): 230-243.
 526
 527 Pöhlmann-Dietze P, Ulrich M, Kiser KB, Döring G, Lee JC, Fournier JM, et al. Adherence of
 528 *Staphylococcus aureus* to endothelial cells: influence of capsular polysaccharide, global
 529 regulator agr, and bacterial growth phase. Infect Immun. 2000;68(9):4865-71.
 530

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

531 Poston SM, Glancey GR, Wyatt JE, Hogan T, Foster TJ. Co-elimination of mec and spa
 532 genes in *Staphylococcus aureus* and the effect of agr and protein A production on bacterial
 533 adherence to cell monolayers. J Med Microbiol. 1993;39(6):422-8.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

534
 535 Wesson CA, Liou LE, Todd KM, Bohach GA, Trumble WR, Bayles KW. *Staphylococcus*
 536 *aureus* Agr and Sar global regulators influence internalization and induction of apoptosis.
 537 Infect Immun. 1998;66(11):5238-43.

538
 539 Ballhausen B, Jung P, Kriegeskorte A, Makgotlho PE, Ruffing U, von Müller L, et al. LA-
 540 MRSA CC398 differ from classical community acquired-MRSA and hospital acquired-
 541 MRSA lineages: functional analysis of infection and colonization processes. Int J Med
 542 Microbiol. 2014;304(7):777-86. doi: 10.1016/j.ijmm.2014.06.006.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

543
 544 Ezepechuk YV, Leung DY, Middleton MH, Bina P, Reiser R, Norris DA. Staphylococcal
 545 toxins and protein A differentially induce cytotoxicity and release of tumor necrosis factor-
 546 alpha from human keratinocytes. J Invest Dermatol. 1996;107(4):603-9.

547
 548 Popov LM, Marceau CD, Starkl PM, Lumb JH, Shah J, Guerrero D, et al. The adherens
 549 junctions control susceptibility to *Staphylococcus aureus* α -toxin. Proc Natl Acad Sci U S A.
 550 2015;112(46):14337-42. doi: 10.1073/pnas.1510265112.

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

551
 552 Grundmeier M, Hussain M, Becker P, Heilmann C, Peters G, Sinha B. Truncation of
 553 fibronectin-binding proteins in *Staphylococcus aureus* strain Newman leads to deficient

554 adherence and host cell invasion due to loss of the cell wall anchor function. Infect Immun.
555 2004;72(12):7155-63.
556
557 Bur S, Preissner KT, Herrmann M, Bischoff M. The *Staphylococcus aureus* extracellular
558 adherence protein promotes bacterial internalization by keratinocytes independent of
559 fibronectin-binding proteins. J Invest Dermatol. 2013;133(8):2004-12. doi:
560 0.1038/jid.2013.87.
561
562

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

Feldfunktion geändert

563 Supporting information

564 **S1 Fig. Phylogenetic tree.** Phylogenetic tree of whole genome sequenced isolates in the
565 context of other sequenced CC398 strains. Scale bar shows the number of substitutions within
566 the expected core genome (1,799,838 bp).

567
568 **S2 Table. Variable elements.** Table providing an overview of variable elements within the
569 genomes of porcine LA-MRSA strains Fin47_H17, Fin46_17, and Fin48_H18 compared to
570 reference strain S0385.

571

572

573

574

575

576

577

578

579

580

581

582 **Supplementary Table 1: Genomic analysis of adhesion associated genes.** S0385 data from Uhlemann 2012, with additions. Abbreviations:
583 V/S = variable/same (compared to strain indicated in superscript), A = absent at this location, I = insertion, X = aa substitution, ψ = pseudogenes,
584 wt = wild type, aa =amino acid, Δ = deletion, SNP = single nucleotide polymorphism.

| Gene | S0385 locus tag ¹ SAPIG | S085 | NM01 | Fin47_H17 | Fin46_H17 | Fin48_H18 | Comment |
|-------------|--|------------------------|-----------------|--------------------|------------------------|-------------------------|-------------------------------|
| <i>spa</i> | 0122 | 8 aa Δ at aa390 | 2 X at aa390 | wt | wt | I (8 aa at aa390) | Variable repeat length |
| <i>coa</i> | 0240 | 81 bp Δ | wt | wt | wt | wt | Variable repeat length |
| <i>vwb</i> | 0483 | wt | A | V ^{S0385} | V ^{S0385} | A (at this location) | Variable sequence/presence |
| <i>sdrC</i> | 0636 | 174 bp Δ | wt | 1 SNP | S ^{Fin47_H17} | I (24aa at aa737) | Variable repeat length |
| <i>sdrD</i> | 0637 | 42 bp Δ and 4 X | Δ 54 bp | wt | wt | wt | Variable repeat length |

| | | | | | | | |
|--------------|--------|-------------------------------|----|--------------------|------------------------|--------------------------|------------------------|
| | | | | | | | and SNPs |
| <i>eno</i> | 0855 | | | S ^{S0385} | S ^{S0385} | 1 SNP, non synonymous | |
| <i>clfA</i> | 0866-7 | ψ truncated, 2 frameshifts | wt | V ^{S0385} | S ^{Fin47_H17} | V ^{S0385} | Variable repeat length |
| <i>isdB</i> | 1125 | SNP, 9 bp Δ | wt | S ^{S0385} | S ^{S0385} | I ^{S0385} | |
| <i>isdA</i> | 1126 | | | S ^{S0385} | S ^{S0385} | 1 SNP, synonymous | |
| <i>ecb</i> | 1150 | wt | wt | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| <i>fib</i> | 1154 | wt | wt | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| <i>ebhA</i> | 1434 | 53 aa Δ at aa9851 and 1 X | wt | 1 aa X | 1 X | 2 X | Variable repeat length |
| <i>ebpS</i> | 1480 | wt | wt | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| hypothetical | 1791 | 174 bp Δ | wt | 1 aa change | S ^{Fin47_H17} | wt | Variable repeat length |

| | | | | | | | |
|-------------|--------|-------------------------|----------------|---------------------------------------|------------------------|-------------------------|------------------------------|
| | | | | (aa39) | | | |
| <i>map</i> | 1981 | | | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| <i>sdrH</i> | 2069 | wt | Δ 39 bp | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| <i>hlD</i> | 2072 | wt | wt | ψ A | wt | wt | Variably present |
| <i>agrB</i> | 2073 | wt | wt | A | wt | wt | Variably present |
| <i>agrA</i> | 2076 | | | S ^{S0385} | S ^{S0385} | S ^{S0385} | |
| <i>fnbB</i> | 2550-1 | ψ truncated, stop codon | wt | Deleted | Deleted | ψ truncated, stop codon | Often defunct |
| <i>fnbA</i> | 2553 | wt | wt | Δ9 bp in repeat region, 12 aa changes | S ^{Fin47_H17} | wt | Variable |
| <i>clfB</i> | 2679 | ψ truncated | Δ42aa at aa819 | wt | wt | wt | Variable repeats / truncated |
| <i>isaB</i> | 2688 | | | S ^{S0385} | S ^{S0385} | 1 SNP, | |

585

| | | | | | | | |
|------------|------|----|-------------------|----------------------|----------|----------------------|------------------------|
| | | | | | | synonymous | |
| <i>cna</i> | 2740 | wt | Δ B domain | Δ 187aas, 4 X | As Fin47 | Δ 187aas, 4 X | Variable in this clade |